



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MORTALITY—FOREIGN AND INSULAR—COUNTRIES AND CITIES
(untabulated).

BRAZIL—Santos.—Two weeks ended September 16, 1911. Population, 85,000. Total number of deaths from all causes 40, including measles 1, typhoid fever 1.

GREAT BRITAIN.—Week ended December 2, 1911.

England and Wales.—The deaths registered in 77 great towns correspond to an annual rate of 15.7 per 1,000 of the population, which is estimated at 16,157,797.

Ireland.—The deaths registered in 21 principal town districts correspond to an annual rate of 18.3 per 1,000 of the population, which is estimated at 1,149,495. The lowest rate was recorded at Galway, viz, 3.9, and the highest at Sligo, viz, 65.4, per 1,000 of the population.

Scotland.—The deaths registered in 8 principal towns correspond to an annual rate of 18 per 1,000 of the population, which is estimated at 1,710,291. The lowest rate was recorded at Perth, viz, 14.5, and the highest at Aberdeen, viz, 20.8, per 1,000 of the population. The total number of deaths from all causes was 589, including diphtheria 15, measles 31, scarlet fever 2, typhoid fever 2.

ITALY—Genoa.—Two weeks ended November 30, 1911. Population, 272,077. Total number of deaths from all causes 162, including diphtheria 3, smallpox 3, tuberculosis 15, typhoid fever 1.

RUSSIA—Riga.—Month of September, 1911. Population, 355,000. Total number of deaths from all causes 592, including diphtheria 12, measles 1, scarlet fever 10, typhoid fever 7, typhus fever 1.

SPAIN—Madrid.—Month of November, 1911. Population, 584,117. Total number of deaths from all causes 987, including diphtheria 28, measles 8, smallpox 1, tuberculosis 144, typhoid fever 14, typhus fever 1.

Malaga.—Month of October, 1911. Population, 135,000. Total number of deaths from all causes 328, including diphtheria 2, measles 2, smallpox 24, tuberculosis 22.

Seville.—Month of November, 1911. Population, 158,235. Total number of deaths from all causes 403, including diphtheria 10, smallpox 3, tuberculosis 55.

By authority of the Secretary of the Treasury:

A. H. GLENNAN,

Acting Surgeon General,

United States Public Health and Marine-Hospital Service.